DANGER ASSESSMENT

Jacquelyn C. Campbell, PhD, RN Copyright © 1985, 1988, 2001

Several risk factors have been associated with homicides (murders) of both batterers and battered women in research conducted after the murders have taken place. We cannot predict what will happen in your case, but we would like you to be aware of the danger of homicide in situations of severe battering and for you to see how may of the risk factors apply to your situation.

Using the calendar, please mark the approximate dates during the past year when you were beaten by your husband or partner. Write on that date how bad the incident was according to the following scale:

- 1. Slapping, pushing; no injuries and/or lasting pain
- 2. Punching, kicking; bruises, cuts, and/or lasting pain
- 3. "Beating up"; severe contusions, burns, broken bones
- 4. Threat to use weapon; head injury, internal injury, permanent injury
- 5. Use of weapon; wounds from weapon

(If **any** of the descriptions for the higher number apply, use the higher number.)

Mark **Yes** or **No** for each of the following. ("He" refers to your husband, partner, ex-husband, ex-husband, or whoever is current physically hurting you.)

1.	Has the physical violence increased in severity or frequency over the past year?
2.	Has he ever used a weapon against you or threatened you with a weapon?
3.	Does he ever try to choke you?
4.	Does he own a gun?
4. 5.	Has he ever forced you to have sex when you did not wish to do so?
6.	Does he use drugs? By drugs, I mean "uppers" pr amphetamines, speed, angel dust, cocaine, "crack", street
	drugs or mixtures.
7.	Does he threaten to kill you and/or do you believe he is capable of killing you?
8.	Is he drunk every day or almost every day? (In terms of quantity of alcohol.)
8. 9.	Does he control most or all of your daily activities? For instance: does he tell you who you can be friends
	with, when you can see your family, how much money you can use, or when you can take the car? (If he
	tries, but you do not let him, check here:)
10.	Have you ever been beaten by him while you were pregnant? (If you have never been pregnant by him,
	check here:)
11.	Is he violently and constantly jealous of you? (For instance, does he say "If I can't have you, no one can.")
12.	Have you ever threatened or tried to commit suicide?
13.	Has he ever threatened or tried to commit suicide?
14.	Does he threaten to harm your children?
15.	Do you have a child that is not his?
16.	Is he unemployed?
17.	Have you left him during the past year? (If you <i>never</i> lived with him, check here:)
	Do you currently have another (different) intimate partner?
19.	Does he follow or spy on you, leave threatening notes, destroy your property, or call you when you don't
	want him to?
	Total "Yes" Answers

Thank you. Please talk to your nurse, advocate or counselor about what the Danger Assessment means in terms of your situation.

Reference List

Block, C.R., Engel, B. Naureckas, S.M., & Riordan, K.A. (1999). The Chicago women's health risk study: Lessons in collaboration. Violence Against Women, 5, 1158-1177.

Campbell, J. (1986). Nursing assessment for risk of homicide with battered women. <u>Advances in Nursing Science</u>, 8, 36-51.

Campbell, J. (1995). Assessing dangerousness. Newbury Park: Sage.

Campbell, J., Soeken, K., McFarlane, J., & Parker, B. (1998). Risk factors for femicide among pregnant and nonpregnant battered women. In J.C. Campbell (Ed.), <u>Empowering survivors of abuse: Health care for battered women and their children</u> (pp. 90-97). Thousand Oaks, CA: Sage.

Diaz-Olavarrieta, C., Campbell, J., Garcia de le Cadena, C., Paz, F., & Villa, A. (1999). Domestic violence against patients with chronic neurologic disorders. <u>Archives of Neurology</u>, 56, 681-685.

McFarlane, J., Parker, B., Soeken, K., & Bullock., L. (1992). Assessing for abuse during pregnancy: Severity and frequency of injuries and associated entry into prenatal care. <u>JAMA</u>, 267, 3176-3178.

McFarlane, J., Parker, B., & Soeken, K. (1996). Abuse during pregnancy: Associations with maternal health and infant birth weight. Nursing Research, 45, 37-42.

McFarlane, J., Soeken, K., Reel, S., Parker, B., & Silva, C. (1997). Resource use by abused women following an intervention program: Associated severity of abuse and reports of abuse ending. <u>Public Health Nursing</u>, 14, 244-250.

Parker, B., McFarlane, J., & Soeken, K. (1994). Abuse during pregnancy: Effects on maternal complications and birth weight in adult and teenage women. <u>Obstetrics & Gynecology</u>, 84, 323-328.

Roehl, J. & Guertin, K. (1998). <u>Current use of dangerousness assessments in sentencing domestic violence offenders</u>. Pacific Grove, CA: State Justice Institute.

Silva, C., McFarlane, J., Soeken, K., Parker, B., & Reel, S. (1997). Symptoms of post-traumatic stress disorder in abused women in a primary care setting. Journal of <u>Women's Health</u>, 6, 543-552.

Stuart, E.P. & Campbell, J.C. (1989). Assessment of patterns of dangerousness with battered women. <u>Issues Mental Health Nursing</u>, 10, 245-260.

Websdale, N. (1999). Understanding domestic homicide. Boston: Northeastern.

Weisz, A., Tolman, R., & Saunders, D.G. Assessing the risk of severe domestic violence: The importance of survivor's predictions. <u>Violence & Victims</u> (in press)

Woods, S. (1998). Post-traumatic stress disorder in battered women. Unpublished doctoral dissertation, Washington State University.